

## Inflection of Beating Transformation by Routing Point on Vitreous Chamber

Jeong-Lae Kim<sup>1</sup>, Eun-Young Kang<sup>2</sup>, Yun-Sik Lim<sup>3</sup>, Jong-Youel Park<sup>4</sup>, Sung-Ho Hwang<sup>5</sup> and Jeong-Jin Kang<sup>6</sup>

### Abstract

*Moist skin on the wild surface is a roll alteration that transports droplets like a superconductor in a single or multi-layer multi-block module. Coruscating-individualization realization level (CIRL) within the skin used a technique of messaging with a roll realization level function in the stiff swell-repercussion status and mixing with pulse changes. In the swell-repercussion status, the roll messaging function within the skin was calculated as the roll value found as a peel-stick-wall-dot by the superstructure being observed. The roll repercussion function shown in the dermis layer consisted of an alteration signal as a realization level based on the coruscating-individualization level, and was measured using the technical concept of pulse change of the diffusion repercussion function and converted to the value of CIRL by the maximum average. The degree of pulse change was expressed as the degree of water distribution within the skin, and a value was formed according to the realization level of the repercussion function. In Ro-RG-FA-ΦMAX-MIN, the roll far alteration value due to roll peel-stick-wall-dot repercussion was shown as  $19.77 \pm 4.36$  units. In roll convenient alteration value, Ro-RG-CO-ΦMAX-MIN was found to be  $6.68 \pm 1.33$ . In roll flank alteration value, Ro-RG-RO-ΦMAX-MIN was found to be  $1.64 \pm 0.01$  units. In roll vicinage alteration value, Ro-RG-VI-ΦMAX-MIN was found to be  $0.42 \pm 0.15$  units. In the dermis layer, swell repercussion by CIRL is evaluated by the degree of peel-stick-wall-dot diffusion function shown at the roll realization level of porous matter, and the dermis layer shown at the messaging level system is expressed as a differentiated function and can be used. You can. The function of the porous in the dermis layer is a concept that suppresses differential signals in diffusion realization systems and could be utilized as a promising key tool to utilize the data from a nanotechnology perspective and for basic applications for analyzing surfaces.*

**Keywords:** Roll Realization Level, Roll Realization Gestalt, Swell Realization System, Swell Repercussion

## INTRODUCTION

The structure of body gestalts that come in contact with a lot of water, such as skin gestalts, is often used for long-term storage of water on the porous surface, hydration, and short-term storage of energy. The extensive scale of structures in porous space is well known to be highly tortuous, disordered, and complex. The complexity of the porous media has a significant impact on its electrical conductivity, permeability and other properties. It is difficult to quantitatively investigate how the complexity of the porous space affects different fibrous tissues. Most other fibrous tissues are composed of crystalline water droplets of various shapes and sizes [1]. Most tissues are composed of epithelial tissue, muscle tissue, nervous tissue, and fibrous tissue. Fibrous tissue is found among all other tissues in the body, including the nervous system. Fibrous tissue is also composed of the connective tissue of the membranes surrounding the brain and spinal cord [2]. Most fibrous tissue types consist of elastic and collagen fibers, ground substance, and cells. Connective tissue is a special type of fluid that does not contain fiber and is classified as blood and lymph. Other connective tissue cells include fibroblasts, adipocytes, macrophages, mast cells, and leukocytes [3]. The human skin tissue consists of the epidermis, dermis, and subcutaneous fat layer. The weight of an adult's skin is approximately 5 kg, and the epidermis is composed of a thin layer, although the thickness varies from 0.04 mm to 1.6 mm depending on the area. The epidermis is a stratified squamous cornified epithelium that goes through several stages of

<sup>1</sup> Department of Biomedical Engineering, Eulji University, Seongnam, 13135, Korea

<sup>2</sup> Department of Computer Software Engineering, Dongyang Mirae University, Seoul, Korea

<sup>3</sup> Department of Electrical Engineering, Yeosu Institute of Technology, 12652 Yeosu, Korea. (Corresponding Author/ E-mail: elecys@yit.ac.kr)

<sup>4</sup> Department of Smart IT, Baewha Women's University, Seoul, Korea

<sup>5</sup> Department of Information and Communication Engineering, Kangwon National University, Samcheok, Korea

<sup>6</sup> Department of Information and Communication, Dong Seoul University, Seongnam, Korea

differentiation of keratinocyte, the main constituent cells, to form a characteristic layer. Fibrous tissue has a wide variety of gestalts that depend on the types of cells and has dense irregular connective tissue and formed mainly by fibroblasts and collagen fibers. Fibrous tissue has an important role in providing a medium for oxygen and nutrients to diffuse from capillaries to cells. This capacity has gestalt of carbon dioxide and waste substances to diffuse from cells back into circulation. They allow organs to resist stretching and tearing forces. They found in highly specialized organs that found in highly specialized organs [4].

Roll repercussion is a sharp tip fastened to the free end of a small one shape that fibrous tissue is rest situation and can be fragments of simple mechanical shapes. The structure of the created epidermis can be divided into the basal layer, spinous layer, granular layer, and stratum corneum, and about 80% of the cells are known to be responsible for keratinocytes, melanocytes that display pigmentation, Langerhans cells that perform immune function, and skin nerves [5]. Merkel cells account for the largest portion. In the epidermis, the barrier function carries out the differentiation of the epidermis, and the most important role in the barrier function is the substances that make up the compartment model of the proteins and lipids of the stratum corneum, which appear as products of epidermal differentiation. The differentiation process starting from epidermal stem cells to differentiated keratinocytes exists with complex regulatory functions. Normal differentiation of the skin is a critical factor in understanding skin barrier function. After the cells divide and proliferate in the basal layer, they move to the upper layer to form the spiny layer, and then to the upper layer to form the granular layer, and finally form the stratum corneum, the outermost layer of the epidermis, which appears on the outside. The layer that appears on the outside protects against bacterial invasion, traction, tension, and mechanical stimulation, and prevents evaporation of moisture, maintaining homeostasis in the human body. The epidermis contains Langerhans cells, which perform immune functions, and melanocytes, which synthesize melanin to protect the body from damage from ultraviolet rays [6]. The roll alteration technique applied to the skin structure roll alterations the coruscating-individualization gestalt to produce stiffen realization of porous matter. Stiffen gestalts distill the roll value of the coruscating-individualization level into the structuralize realization of the individualization peel-stick-wall-dot. Roll alteration causes the peel-stick-wall-dot to distill the roll value by structuralizing the upper swell. Swell-repercussion becomes a peel-stick-wall-dot depending on the level of roll realization and mixes the capabilities of the swell gestalt. The porous surface of moisture by roll realization gestalts system seeks to recognize the coruscating-individualization realization level.

## THEORY

### Roll Realization Messaging

Roll realization gestalt (Ro-RG) is wild messaging to define immixture valued with upper layer peel-stick-wall-dot on the repercussion. The porous matter is a roll realization gestalt that structuralizes realization that Ro-RG is Overall Repercussion Level (OSL), Far-Convenient Repercussion Level (FCEL) and Flank-Vicinage Repercussion Level (FVEL). Ro-RG levels are standard deviations to investigate with path of phase vicinage side layer from main-peel-stick-wall-dot. The skin structure causes stiffen realization of porous matter that Ro-RG levels are to immixture in degrees. Ro-RG repercussion level scores receive the integrate dislocation for stiffened structuralize signal by swell peel-stick-wall-dot gestalt that causes stiffen realization of far-convenient (FC) and flank-vicinage (FV). Ro-RG dislocation is from horizontal along Ro-FC-axes at x-direction. Vertical along Ro-FV-axes is at y-direction. Ro-FC-axes and Ro-FV-axes were investigated by Ro-RG-FC and Ro-RG-FV respectively. FVEL can immixture of the received structuralize signal both amplitude and phase. I-Ro-RG and Q-Ro-CV applied to the skin structure, is to current far-convenient and flank-vicinage by the Ro-RG-FV and Ro-RG-FC. Ro-FC is roll alteration technique applied to the skin structure on Ro-RG modulated carrier of roll far-convenient. Ro-FV is roll alteration technique applied to the skin structure on Ro-RG modulated carrier of roll flank-vicinage.  $\Delta P_{Ro-RG}$  is a on the Ro-RG mplitude and phase of the received structuralize signal of the roll  $I_{Ro-FC}$  and roll  $Q_{Ro-FV}$  [7,8](1,2). Equation (1,2) on the absolute value  $\Delta_\gamma$  is investigated as the  $\Delta P_{Ro-RG-FC}$  and  $\Delta P_{Ro-RG-FV}$ .

$$\Delta P_{Ro-KF} = \frac{I_{Ro-FC}^2 + Q_{Ro-FV}^2}{Z_0}, \varphi = \arctan \frac{Q_{Ro-FV}}{I_{Ro-FC}} \quad (1)$$

$$|\Delta_\gamma| = \sqrt{I_{\text{Ro-FC}}^2 + Q_{\text{Ro-FV}}^2} = \sqrt{\Delta P_{\text{Ro-FV-FC}} + Z_0} \quad (2)$$

Indirectly immixture upper layer peel-stick-wall-dot score data, to denote as  $\Delta_\gamma$ , is concerned to the differential reflection coefficient Ro-RG-FC and Ro-RG-FV, can found as (3).  $Z_0$  is received input impedance.

$$\angle(\Delta_\gamma) = \arctan \frac{Q_{\text{Fl-FV}}}{I_{\text{Fl-FC}}} = \phi \quad (3)$$

Roll realization gestalt that structuralizes realization is included by examination setting. The communication range between roll layer pin and their system to comprise of the properly adhere by monitoring [9].

### **Swell Upper Layer Gestalt (Swe-ULG)**

Swell upper layer gestalt (Swe-ULG) requires a combination scores to swell peel-stick-wall-dot gestalt that causes stiffen realization by Swe-ULG-FV and Swe-ULG-FC. Swe-ULG-value is counts to roll realization gestalt is applied absolute  $\Omega$ -Ro-RG values. FV-FC and  $\Omega$ -Ro-RG level alterations are more sensitivity Swe-ULG recovered to employ wide individualization propagation shape (4) to  $\Omega$ -Ro-RG based on Swe-ULG-FC and Swe-ULG-FV:

$$\Omega\text{-Ro-RG}(r)[\text{n.u.}] = \Omega_{\text{Swe-ULG-FC}} \Omega / r^{\Omega\text{-Swe-ULG-FV}} \equiv \Omega\text{-Ro-RG}(r)[\text{dB}] = 20\log_{10}(\Omega_{\text{Swe-ULG-FV}}) - \Omega_{\text{Swe-ULG-FC}} 20\log_{10}(r) \quad (4)$$

Swe-ULG of 'r' is the range or distance.  $\Omega_{\text{Swe-ULG-FV}}$  and  $\Omega_{\text{Swe-ULG-FC}}$  are coefficients by investigated from a non-multi regression to minimize the root mean square (RMS) on set of between main-peel-stick-wall-dot and side-peel-stick-wall-dot. The expression rate of  $\Omega\text{-Ro-RG}(r)$  is already multi with regard to  $\Omega_{\text{Swe-ULG-FV}}$  and  $\Omega_{\text{Swe-ULG-FC}}$  [10,11].

### **Tolling Perception Figuration Selection**

Supported striking peculiarity of roses-butterflies dot figuration is porous matter is a swell peel-stick-wall-dot gestalt that causes stiffen realization is roses-butterflies dot. Tolling perception figuration (Tol-PF) is the unruly structuralized on the upper layer roses-butterflies dot activity to tie-up sparkle-divergence upper layer level (SDULL) (Figure 1)

### **Roll Realization Gestalt (Ro-RG)**

Roll realization gestalt (Ro-RG) is to incur striking oddity of peel-stick-wall-dot gestalt on dot peel-stick-wall-dot. Stiffened Upper layer peel-stick-wall-dot activity is integrated of structuralized at coruscating-individualization upper layer level (BDULL) (Figure 1). Swell-repercussion peel-stick-wall-dot level (Swe-PSDL) is result to influence on parameter of BDULL. Roll repercussion gestalt (Ro-SF) in coruscating-individualization activity is structuralized of roll repercussion to exercise structuralize [12].

### **Swell Peel-Stick-Wall-Dot Gestalt (Swe-PSDG)**

Ro-RG system is recovered stiffened form in structuralizes realization on roll realization gestalt system (Ro-RGS) as peel-stick-wall-dot. Ro-RG is denoted to look on stiffened swell level. Denote upper layer peel-stick-wall-dot techniques (ULFCT) Ro-RG is to similar to hold down swell-repercussion. Stiffened swell-repercussion is to hold down to integrate in swell upper layer peel-stick-wall-dot gestalt (Swe-PSDG). Stiffened swell-repercussion on peel-stick-wall-dot is propelled tool with roll layer (Ro-L). Arithmetic striking oddity at the roll structuralized (Ro-S) is propelled in swell peel-stick-wall-dot gestalt (Swe-PSDG) to immixture Ro-RGS of output parameters for peel-stick-wall-dot. Swell-repercussion gestalt (Swe-SF) in the Ro-RGS is to recover by Ro-RG on swell realization level (Swe-CL) with immixture of output parameters. Ro-SF on ULFCT of Ro-RG was investigated to upper of layer (UOL) to upper layer swell-repercussion techniques (Swe-RT) of vicinage direction. Roll alteration technique applied to the skin structure causes stiffen realization is swell realization level gestalt (Swe-RLF) on ULFCT of Ro-RG to distill swell signal to layer structuralize mechanisms. Roll coruscating-individualization level (Ro-CIL) on Swe-RLF is found by stiffen realization to swell realization and swell gestalt. Swe-RLF is denote with swell realization gestalt (Swe-RG) of soft swell signal [13] (Figure 2).

**EXPERIMENT**

**Roll realization gestalt (Ro-RG)**

Roll realization gestalt (Ro-RG) on dot peel-stick-wall-dot is roll realization gestalt that structuralizes realization to incur striking oddity of peel-stick-wall-dot gestalt. Upper layer peel-stick-wall-dot activity is integrated stiffened structuralized to coruscating-individualization upper layer level (CIULL) (Figure 1). Swell-repercussion peel-stick-wall-dot level (Swe-SDL) resulted to influence parameter of CIULL. Roll repercussion gestalt (Ro-SF) in coruscating-individualization activity is to exercise of roll repercussion structuralize [14].

**Swell Peel-Stick-Wall-Dot Gestalt (Swe-PSDG)**

Ro-RG system on roll realization gestalt system (Ro-RGS) is to recover stiffened form by peel-stick-wall-dot. Ro-RG of upper layer peel-stick-wall-dot techniques (ULFCT) is denote to recover stiffened swell level is similar to hold down swell-repercussion. Integrated in swell upper layer peel-stick-wall-dot gestalt (Swe-ULGCF) hold down to stiffened swell-repercussion is propelled by roll layer (Ro-L) tool on peel-stick-wall-dot. Arithmetic striking oddity by Ro-RGS is propelled at roll alteration technique applied to the skin structure for peel-stick-wall-dot. The roll structuralized (Ro-S) propel for immixture of output parameters by in swell peel-stick-wall-dot gestalt (Swe-PSDG). Swell-repercussion gestalt (Swe-SF) by Ro-RG is to recover with immixture of output parameters by swell realization level (Swe-CL) in the Ro-RGS. Ro-SF on ULFCT of Ro-RG is investigated for upper of layer (UOL) by upper layer swell-repercussion techniques (Swe-ST) of vicinage direction. Swell realization level gestalt (Swe-CLF) on ULFCT of Ro-RG is distilled by swell signal to layer structuralize mechanisms. Roll coruscating-individualization level (Ro-CIL) on Swe-CLF is found to swell realization and the swell gestalt. Swe-CLF by swell realization gestalt (Swe-CF) is denote to apply for the skin structure on soft swell signal [15] (Figure 2).




Fig. 1. Coruscating-individualization gestalt structuralized roll realization location on the matter.




Fig. 1. Sread-out realization gestalt is system block by coruscating-individualization level on the roll alteration technique.

## RESULTS AND DISCUSSION

### Properties of the Sequence Selection

Ro-RG-gestalt of stiffen realization of porous matter is to examination to define  $Ro-RG-\Phi_{MAX-MIN}$ ,  $Ro-RG-\Phi_{MAX-MED}$  and  $Ro-RG-\Phi_{MAX-AVG}$  database. Ro-RG-gestalt by Ro-RG activities create to accumulate by roll oddity repercussion gestalt (Ro-CRF) (Table 1). Roll repercussion gestalt data for calculations are to employ Matlab6.1.

Table 1. Roll dot gestalt (Ro-DF) average: far RO-BDCL ( $Ro-RG-FA\Phi_{MAX-AVG}$ ), convenient RO-BDCL ( $Ro-RG-CO\Phi_{MAX-AVG}$ ), flank RO-BDCL ( $Ro-RG-FL\Phi_{MAX-AVG}$ ) and vicinage RO-BDCL ( $Ro-RG-VI\Phi_{MAX-AVG}$ ) condition.  $Ro-RG-\Phi_{MAX-AVG}$  and  $Ro-RG-\Phi_{MAX-MIN}$  average.

Average $\Phi$	FA $\Phi_{Avg-RO-BDCL}$	CO $\Phi_{Avg-RO-BDCL}$	FL $\Phi_{Avg-RO-BDCL}$	VI $\Phi_{Avg-RO-BDCL}$
$Ro-RG-\Phi_{MAX-MED}$	$14.13 \pm 5.81$	$4.06 \pm 1.82$	$0.83 \pm 0.19$	$0.28 \pm 0.15$
$Ro-RG-\Phi_{MAX-AVG}$	$11.32 \pm (-2.18)$	$3.56 \pm (-0.72)$	$0.81 \pm (-0.34)$	$0.23 \pm (-0.01)$

### Improvements Of Multiple Alignments by Sequence Selections

Roll realization gestalt (Ro-RG) on repercussion technique (ST) condition is repercussion status to make-out of coruscating-individualization level (CIL). ET of structuralizes realization look for roll coruscating-individualization level (Ro-CIL) on stiffened objects of Ro-RG-gestalt. RT on Ro-RG-gestalt adhere the equivalent things of peel-stick-wall-dot. Roll realization gestalt system (Ro-RGS) of coruscating-individualization realization level (BDCL) is result for oddity in accordance to make-out parameter. BDCL in swell realization gestalt activities (Swe-RGA) is propelled coruscating alteration to denote examination.

### Ro-BDCL of Comparison Database on $Ro-RG-\Phi_{AVG}$ and $Ro-RG-\Phi_{MAX-MIN}$ and $Ro-RG-\Phi_{MAX-MED}$

Roll Roll realization gestalt (Ro-RG) on far (FA- $\Phi$ ) condition is to denote stiffened a roll coruscating-individualization realization level (Ro-BDCL) value;  $Ro-RG-FA-\Phi_{MAX-MIN}$ ,  $Ro-RG-FA-\Phi_{AVG}$  and  $Ro-RG-FA-\Phi_{MAX-MED}$  (Figure 3). Applied the skin structure to causes stiffen realization of porous matter  $Ro-RG-FA-\Phi_{AVG}$  is large roll in Ro-RGS of dot-flank-vicinage (DFV) direction. Far Ro-BDCL in Ro-RGS is structuralizes realization to make-out Ro-RG activities of differential with same direction between  $Ro-RG-FA-\Phi_{AVG}$  and  $Ro-RG-FA-\Phi_{MAX-MED}$ . Realized individualization peel-stick-wall-dot of  $Ro-RG-FA-\Phi_{AVG}$  of roll dot gestalt (Ro-DF) with far Ro-BDCL is structuralizes realization to make-out very large roll Ro-RG activities at  $16.97 \pm 9.24$  unit.  $Ro-RG-FA-\Phi_{MAX-MIN}$  in the Ro-RGS with far Ro-BDCL is structuralizes realization to make-out some

large Ro-RG activities at  $19.77 \pm 4.36$  unit. Ro-RG-FA- $\Phi_{\text{MAX-MED}}$  with far Ro-BDCL on roll coruscating-individualization level is structuralizes realization to make-out some large Ro-RG activities at  $14.13 \pm 5.81$  unit.


Roll dot gestalt (Ro-DF) by far Ro-BDCL in roll activities of Ro-RG-Far is find roll influence to occur the flank-vicinage (FV) direction of far repercussion. Roll realization gestalt (Ro-RG) of convenient (CO- $\Phi$ ) condition is to denote stiffened a roll coruscating-individualization realization level (Ro-BDCL) value; Ro-RG-CO- $\Phi_{\text{AVG}}$ , Ro-RG-CO- $\Phi_{\text{AVG}}$  and Ro-RG-CO- $\Phi_{\text{MAX-MED}}$  (Figure 3). Applied the skin structure to causes stiffen realization of porous matter on convenient Ro-BDCL in Ro-RGS is structuralizes realization to make-out Ro-RG activities of differential with same direction between Ro-RG-CO- $\Phi_{\text{AVG}}$  and Ro-RG-CO- $\Phi_{\text{AVG}}$ . Besides, convenient Ro-BDCL is structuralizes realization to make-out Ro-RG activities of a small roll at Ro-RG-CO- $\Phi_{\text{MAX-MED}}$  of the roll dot gestalt (Ro-DF) on the FV direction in the Ro-RGS. Realized individualization peel-stick-wall-dot of Ro-RG-CO- $\Phi_{\text{AVG}}$  of roll dot gestalt (Ro-DF) with convenient Ro-BDCL is structuralizes realization to make-out some large roll Ro-RG activities at  $8.66 \pm 3.08$  unit. Ro-RG-CO- $\Phi_{\text{MAX-MIN}}$  in the Ro-RGS with convenient Ro-BDCL is structuralizes realization to make-out large of Ro-RG activities at  $6.68 \pm 1.33$  unit. It is a minute role in the roll activities of a convenient repercussion. Ro-RG-CO- $\Phi_{\text{MAX-MED}}$  at roll coruscating-individualization level on the FC direction is structuralizes realization to make-out small roll Ro-RG activities at  $4.06 \pm 1.82$  unit. Roll dot gestalt (Ro-DF) by convenient Ro-BDCL is found a roll at the same direction to occur in the Ro-RGS activities direction.

Roll realization gestalt (Ro-RG) of flank (RO- $\Phi$ ) condition is to denote stiffened a roll coruscating-individualization realization level (Ro-BDCL) value; Ro-RG-RO- $\Phi_{\text{AVG}}$ , Ro-RG-RO- $\Phi_{\text{AVG}}$  and Ro-RG-RO- $\Phi_{\text{MAX-MED}}$  (Figure 3). Applied the skin structure to causes stiffen realization of porous matter on flank Ro-BDCL is structuralizes realization to make-out with small roll on FV direction of Ro-RGS of the roll dot gestalt (Ro-DF) at Ro-RG-RO- $\Phi_{\text{AVG}}$  and Ro-RG-RO- $\Phi_{\text{AVG}}$ . Roll value of Ro-RG-RO- $\Phi_{\text{MAX-MED}}$  is differently the very small to FV direction in Ro-RGS. Realized individualization peel-stick-wall-dot of Ro-RG-RO- $\Phi_{\text{AVG}}$  of roll dot gestalt (Ro-DF) with flank Ro-BDCL is structuralizes realization to make-out small roll Ro-RG activities at  $2.80 \pm 0.74$  unit. Ro-RG-RO- $\Phi_{\text{MAX-MIN}}$  on the FC direction at roll coruscating-individualization level with flank Ro-BDCL is structuralizes realization to make-out small Ro-RG activities at  $1.64 \pm 0.01$  unit. Ro-RG-RO- $\Phi_{\text{MAX-MED}}$  with flank Ro-BDCL is structuralizes realization to make-out very small roll Ro-RG activities at  $0.83 \pm 0.19$  unit.

Ro-BDCL of roll dot gestalt (Ro-DF) in the Ro-RGS activities is to find the same direction to occur a roll of a flank repercussion. Roll realization gestalt (Ro-RG) of vicinage (VI- $\Phi$ ) condition is to denote stiffened a roll coruscating-individualization realization level (Ro-BDCL) value; the Ro-RG-VI- $\Phi_{\text{AVG}}$ , Ro-RG-VI- $\Phi_{\text{AVG}}$  and Ro-RG-VI- $\Phi_{\text{MAX-MED}}$  (Figure 3). Applied the skin structure to causes stiffen realization of porous matter on Ro-RG activities of vicinage Ro-BDCL is structuralizes realization to make-out small roll on FC direction of Ro-RGS at Ro-RG-VI- $\Phi_{\text{AVG}}$  and Ro-RG-VI- $\Phi_{\text{MAX-MIN}}$  of the roll dot gestalt (Ro-DF). Realized individualization peel-stick-wall-dot of Roll value of Ro-RG-VI- $\Phi_{\text{MAX-MED}}$  is differently the small to FV direction in Ro-RGS. Ro-RG-VI- $\Phi_{\text{AVG}}$  of roll dot gestalt (Ro-DF) on roll coruscating-individualization level with vicinage Ro-BDCL is structuralizes realization to make-out very small Ro-RG activities at  $0.55 \pm 0.21$ . Ro-RG-VI- $\Phi_{\text{MAX-MIN}}$  on the FC direction with vicinage Ro-BDCL is structuralizes realization to make-out very small Ro-RG activities at  $0.42 \pm 0.15$  unit. Ro-RG-VI- $\Phi_{\text{MAX-MED}}$  on the FC direction in the Ro-RGS is structuralizes realization to make-out very little small roll Ro-RG activities at  $0.28 \pm 0.15$  unit. Vicinage Ro-BDCL in the roll activities of a vicinage repercussion is to found Ro-RG-VI- $\Phi_{\text{MAX-MED}}$  on the FC direction in the Ro-RGS of roll to occur the same direction.


*Inflection of Beating Transformation by Routing Point on Vitreous Chamber*

**Database on Ro-RG-YOAVG and Ro-RG-YOMAX-MIN and Ro-RG-YOMAX-MED**  
 Roll realization gestalt (Ro-RG) is wild messaging to define immixture valued with upper layer peel-stick-wall-dot on the repercussion. The porous matter is a roll realization gestalt that structuralizes realization that Ro-RG is Overall Repercussion Level (OSL), Far-Convenient Repercussion Level (FCEL) and Flank-Vicinage Repercussion Level (FVEL). Supported striking peculiarity of roses-butterflies dot figuration is porous matter is a swell peel-stick-wall-dot gestalt that causes stiffen realization is roses-butterflies dot. Tolling perception figuration (Tol-PF) is the unruly structuralized on the upper layer roses-butterflies dot activity to tie-up sparkle-divergence upper layer level (SDULL)



Applied the skin structure to causes stiffen realization of porous matter Ro-RG-FA-YOAVG is large roll in Ro-RGS of dot-flank-vicinage (DFV) direction. Far Ro-BDCL in Ro-RGS is structuralizes realization to make-out Ro-RG activities of differential with same direction between Ro-RG-FA-YOAVG and Ro-RG-FA-YOMAX-MED.

**Database on Ro-RG-YOAVG and Ro-RG-YOMAX-MIN and Ro-RG-YOMAX-MED**  
 Ro-RG-gestalt of stiffen realization of porous matter is to examination to define Ro-RG-YOMAX-MIN, Ro-RG-YO-MAX-MED and Ro-RG-YOMAX-AVG database. Ro-RG-gestalt by Ro-RG activities create to accumulate by roll oddity repercussion gestalt (Ro-CRF). Roll realization gestalt (Ro-RG) on repercussion technique (ST) condition is repercussion status to make-out of coruscating-individualization level (CIL). ET of structuralizes realization look for roll coruscating-individualization level (Ro-CIL) on stiffened objects of Ro-RG-gestalt. RT on Ro-RG-gestalt adhere the equivalent things of peel-stick-wall-dot..



Applied the skin structure to causes stiffen realization of porous matter Ro-RG-FA-YOAVG is large roll in Ro-RGS of dot-flank-vicinage (DFV) direction. Far Ro-BDCL in Ro-RGS is structuralizes realization to make-out Ro-RG activities of differential with same direction between Ro-RG-FA-YOAVG and Ro-RG-FA-YOMAX-MED.




Fig. 3. Ro-RG-gestalt of the data on the roll condition for activities: parameter of the  $Ro-RG-\Phi_{AVG}$  and  $Ro-RG-\Phi_{MAX-AVG}$ .

## CONCLUSION

This paper considered various roll repercussion functions (Ro-RF) to determine a repercussion realization suitable for the coruscating-individualization realization level (CIRL), provided a separate swell alteration technique that can be used for the roll realization function, and presented the performance. The realization rate expressed in the value of the roll repercussion function (Ro-RF) was used as the basic reference value of the coruscating-individualization level (CIL) to obtain transition data. The roll values selected for coruscating peel-stick-wall-dot were performed so that they could be used in the same empirical study, and a new evaluation was made using the proposed method and performance with the differential function obtained from the roll realization level system in swell repercussion. In basic applications and nanotechnology, the diffusion realization system received from the differential signal can be used as porous matter data to suppress and analyze the function of the skin structure.

## REFERENCES

- Biga, Lindsay M.; Dawson, Sierra; Harwell, Amy; Hopkins, Robin; Kaufmann, Joel; LeMaster, Mike; Matern, Philip; Morrison-Graham, Katie; Quick, Devon (2019), "4.3 Connective Tissue Supports and Protects", Anatomy & Physiology, OpenStax/Oregon State University, retrieved 16 April 2021
- Fluid Tissues". Biology LibreTexts. 21 May 2021. Retrieved 2 August 2022.
- Mathews, M. B. (1975). Connective Tissue, Macromolecular Structure Evolution. Springer-Verlag, Berlin and New York. link.
- Ross M, Pawlina W (2011). Histology: A Text and Atlas (6th ed.). Lippincott Williams & Wilkins. pp. 158–97. ISBN 978-0781772006.
- Young B, Woodford P, O'Dowd G (2013). Wheater's Gestaltal Histology: A Text and Colour Atlas (6th ed.). Elsevier. p. 65. ISBN 978-0702047473
- Soumya Ranjan Nayak a, JibiteshMishra b, Gopinath Palai, Analysing roughness of surface through subject dimension: A review, Image and Vision Computing, 89 (2019) 21–34.
- Huiting J, Flisijn H, Kokkeler ABJ, Smit GJM. Exploiting phase measurements of EPC Gen2 RFID structures. IEEE Int Conf RFID-Technol Appl (RFID-TA), (2013), 1–6.
- Bekkali A, Zou SC, Kadri A, Crisp M, Penty RV., Performance analysis of passive UHF RFID systems under cascaded fading channels and interference effects. IEEE Trans Wirel Commun., 14 (3) (2015), 1421–33.
- DiGiampaolo E, Martinelli F, Mobile robot localization using the phase of passive UHF RFID signals. IEEE Trans Ind Electron, 61(1) (2014), 365–76.



*Inflection of Beating Transformation by Routing Point on Vitreous Chamber*

- López Y. Á., Gómez M.E., Andrés F.L.H., A received signal strength RFID-based indoor point system, *Sensors and Actuators A*, 255 (2017), 118–133.
- Chawla K., McFarland C., Robins G., Shope C., Real-time RFID localization using RSS, in: 2013 International Conference on Localization and GNSS (ICL-GNSS), Turin (Italy), (2013)(25–27 June), 1–6.
- V301-1,302-1,306-p// Kim, J.L., Choi, J.S. and Hwang, K.S., A Study on Anticipation System of Shudder Distinction by the Physical Shape Alteration in Static Condition, *The Journal of IIBC (JIIBC)*, 17(3) (2017), pp.115-120, DOI 10.7236/JIIBC.2017.17.3.115
- Kim, J.L. and K.D. Kim, Prediction of shiver differentiation by the form alteration on the stable condition, *International Journal of Internet Broadcasting and Communication (IJIBC)*, 9(4) (2017), 8-13. DOI 10.7236/IJIBC.2017.9.4.8
- [V308] ] ] Kim, J.L. and Hwang, K.S., Study of rough wavelength of dynamic alteration with posture, *International Journal of Advanced Smart Convergence (IJASC)*, 4(1) (2015), 99-103.
- Kim, J.L. and Kim, K.D., Denotation of central motion techniques: limpness motion gestalt and limpness sensory unit gestalt, *International Journal of Advanced Culture Technology (IJACT)*, 4(3) (2016), 56-61. DOI 10.17703/IJACT.2016.4.3.56